The Antidote Formulation Instructions			
Step 1	**Fill your Acidification Container with:** **10 drops Part A, plus 10 drops Part B = 1ml** *(Part A can be 20 to 25% NaClO2, Part B is 4% HCl)*		
Step 2	**Stir & wait 5 minutes for CLO2 generation** **(Keep the container standing straight up)**		
Step 3	**After 5 minutes add above CLO2 mixture to one volume of water below** ⬇	⬇	⬇
Bottled Water	**100ml Water**	**500ml Water**	**1000ml Water**
Final Concentration	300 ppm	60 ppm	30 ppm
Step 4	**Label your bottle with the ppm & use right away. Refrigerate if possible, & keep out of sunlight.**		

The Antidote Concentration Usage Instructions for Viral, Bacterial & Fungal Infections plus Cancer	
300 ppm	Use as a powerful disinfectant/sanitizer solution for hands, wounds, surfaces, packages, etc. (Not for eyes)
60 ppm	Use as nebulizing mist for throat, lungs and nose. As a gargle, a mouthwash, for brushing teeth. **Doctors: The Antidote** can be used in an IV for viral, bacterial & fungal infections, cytokine storm, sepsis & cancer. (250-1000cc, 14 drops/min (aprx. 1000cc per 24 hours). *Note: can increase up to 150 ppm in 1000cc, 14 drops/min, for up to 3 days in severe cases.
30 ppm	**Viral, Bacterial & Fungal infections + Cancer (oral use):** make 1 liter of 30ppm & drink 100ml every 30 min until finished. Pause 2 hours & then repeat with another liter. If very sick, make 1 liter & drink 100ml every 15 min until finished. Pause 1 hour and then repeat. Then pause 2 hours & switch to 100ml every 30 min with 3rd liter. Increases blood oxygen saturation, kills bacteria & fungal infections. Neutralizes viruses & toxins via oxidation. Helps to detoxify the body which assists the immune system. **Post Cov-19 patients & Prevention: Prevention:** Drink 1 liter/day at rate of 100ml per hour for at least 14 days after contact with infected patient. **Post Cov-19 patients:** Same as prevention but for 30 days to reduce chance of post Cov-19 complications
Important Please Read ⇨	**While taking The Antidote:** Do not take vitamin C, ascorbic acid or antioxidants (may take 2 hours after stopping **The Antidote** at end of day). Avoid flour, sugars carbohydrates & dairy. Pause drinking for 1 hour after meals. Before & after medication stop for 1 hour. Stop for 30 min after coffee or tea. ***Cov-19 Patients:** Consider fasting & only drinking water while sick. Keep all concentrations out of direct sunlight. No need to refrigerate if used within a few days but refrigerate whenever possible. Tastes better cold. Drink from bottle with small mouth to avoid smelling. 4 to 5 swallows are about 100ml. All dosing here is for average adult – reduce volume of dose proportionally for children & increase for large person but keep the same ppm concentrations. ***Note:** **Use The Antidote immediately**. If concentrations are made & then immediately refrigerated & not used, they can gradually increase slightly as follows: 300 ppm → 400 ppm | 60 ppm → 80 ppm | 30 ppm → 40 ppm. Warning! DO NOT INHALE PURE CLO2 GAS. DO NOT allow undiluted concentration to get into eyes or on skin. Rinse with water for 15 min. Seek medical help for eyes.

The Antidote Acidification Container Examples

Below are examples of possible "Acidification Containers" to give some ideas of what can be used. The important thing is that the diameter isn't too big, ideally a little smaller than your little finger or up to just a little bigger.